

ZASSENHAUS CONJECTURE FOR A_6

MARTIN HERTWECK

Dedicated to the memory of I. S. Luthar[†]

ABSTRACT. For the alternating group A_6 of degree 6, Zassenhaus' conjecture about rational conjugacy of torsion units in integral group rings is confirmed.

1. INTRODUCTION

It has proven exceedingly difficult to achieve progress in a number of areas revolving around units in integral group rings of finite groups, and this applies in particular to the conjecture of the title:

(ZC1) For a finite group G , every torsion unit in its integral group ring $\mathbb{Z}G$ is conjugate to an element of $\pm G$ in the units of the rational group algebra $\mathbb{Q}G$.

It remains not only unsolved but also lacking in plausible means of either proof or counter-example. Only a few non-solvable groups G are known to satisfy (ZC1); in this note, the conjecture is verified for the alternating group A_6 . In 1988, Luthar and Passi [17] started investigations in this field when dealing with A_5 , thereby introducing what is now called the “Luthar–Passi method.” A little later, Luthar and Trama [20] established (ZC1) for the symmetric group S_5 , with a large part of the calculations being done inside the integral group ring $\mathbb{Z}S_5$, or rather a Wedderburn embedding, itself. Mentioning further that the covering group of A_5 can be dealt with some additional arguments [9], that accurately describes the state of knowledge at the turn of the century. The limitations of the original version of the Luthar–Passi method were shown in [2]. Recently, a modular version, especially suited for application to non-solvable groups, was given in [13], and used to verify (ZC1) for the groups $\mathrm{PSL}(2, p)$, $p = 7, 11, 13$. (At that point, the reader may wish to recall that $A_6 \cong \mathrm{PSL}(2, 9)$.) A brief review of the method is given in Section 2. It was also used in [7] to verify (ZC1) for a covering group of S_5 and for $\mathrm{GL}(2, 5)$. The method makes use of the (ordinary) character table and of modular character tables, in a process suited for being done on a computer, the result being that rational conjugacy of torsion units of a given order to group elements is either proven or not, and if not, at least some information about partial augmentations is obtained (see [3–6]).

It should be remarked that from the very beginning, investigations on (ZC1) concentrated on solvable groups. Here we only mention that Luthar, in collaboration

Date: February 2, 2008.

2000 Mathematics Subject Classification. Primary 16S34, 16U60; Secondary 20C05.

Subject classification. RT.

Key words and phrases. integral group ring, torsion unit, Zassenhaus conjecture.

[†]Professor Luthar died at the age of 74 in December 2006.

with others, initiated the study of metacyclic groups [1, 16, 18, 19]. For the whole class of metacyclic groups, (ZC1) has been confirmed only very recently in [12]. Other, more recent, results can be found in [14] and [15].

We briefly recall some of the necessary background. Let G be a finite group. For a group ring element $u = \sum_{g \in G} a_g g$ (all a_g in $\mathbb{Z}$), its partial augmentation $\varepsilon_x(u)$ with respect to an element x of G , or rather its conjugacy class x^G in G , is the sum $\sum_{g \in x^G} a_g$. The augmentation of u is the sum $\sum_{g \in G} a_g$ of all of its partial augmentations. Of course, we need to consider only units of augmentation one in $\mathbb{Z}G$, which form a group we denote by $V(\mathbb{Z}G)$.

A criterion for a torsion unit u in $V(\mathbb{Z}G)$ to be conjugate to an element of G in the units of $\mathbb{Q}G$, which is especially suited for computational purposes, is that all but one of the partial augmentations of every power of u vanish (see [21, Theorem 2.5]). The partial augmentations of u one has to turn attention to are limited by the following remark.

Remark 1. Let u be a torsion unit in $V(\mathbb{Z}G)$. Then $g \in G$ and $\varepsilon_g(u) \neq 0$ implies that the order of g divides the order of u . Indeed, it is well-known that then prime divisors of the order of g divide the order of u (see [21, Theorem 2.7], as well as [14, Lemma 2.8] for an alternative proof). Further, it was observed in [13, Proposition 2.2] that the orders of the p -parts of g cannot exceed those of u .

It is known (see [8]) that the order of a torsion unit u in $V(\mathbb{Z}G)$ is a divisor of the exponent of G . However, it is not known whether some element of G has the same order as u . For example, the more interesting part we will be concerned with when proving (ZC1) for A_6 is about the non-existence of torsion units of order 6 in $V(\mathbb{Z}A_6)$, despite the fact that there are no elements of order 6 in A_6 (see Section 4). Using the Luthar–Passi method alone is not sufficient. In this situation, we are lucky in that only in a rough sense ‘ties’ between two integral representations (affording the 5-dimensional irreducible characters) need to be considered.

In Section 3, we apply the Luthar–Passi method to torsion units in $V(\mathbb{Z}A_6)$, with the result that all torsion units not of order 6 are conjugate to elements of A_6 in the units of $\mathbb{Q}A_6$, and if there should exist a unit of order 6, its partial augmentations are essentially unique and can be specified. That is a matter of routine, and was already done by Salim [22] who calculated the coefficients of the linear inequalities arising from equation (2) below, and then their solutions. We shall proceed in a straightforward way, making only use of inequalities arising from equation (1).

2. THE LUTHAR–PASSI METHOD

We briefly survey, in an informal way, the Luthar–Passi method. Still, let u be a torsion unit in $V(\mathbb{Z}G)$, of order n (say). Then, conjecture (ZC1) states that there is $g \in G$ such that $\chi(u^i) = \chi(g^i)$ for all irreducible characters χ of G and integers i . This is because complex representations are determined (up to equivalence) by their characters, and conjugacy in the units of $\mathbb{C}G$ can be shown to be equivalent to conjugacy in the units of $\mathbb{Q}G$ (“rational conjugacy”). It is perhaps even more natural to take the eigenvalues of representing matrices into consideration: If an irreducible character χ is afforded by a complex representation D of G , write $\mu(\xi, u, \chi)$ for the multiplicity of an n -th root of unity ξ as an eigenvalue of the matrix $D(u)$. Then u is rationally conjugate to the group element g if and only if $\mu(\xi, u, \chi) = \mu(\xi, g, \chi)$ for all χ and ξ .

The same applies, *mutatis mutandis*, to Brauer characters. Suppose that u is a p -regular torsion unit for some prime p dividing the order of G (that is, p does not divide the order n of u). A Brauer character φ (relative to p) is a complex-valued function associated with a modular representation of G (in characteristic p), defined on the set of p -regular elements of G . The way Brauer defined these functions actually shows that the domain of φ can be extended to the set of p -regular torsion units of $V(\mathbb{Z}G)$, and the decomposition matrix gives the relation between the values of irreducible Brauer characters and ordinary characters on these units. Multiplicities $\mu(\xi, u, \varphi)$ are defined in the obvious way. Then one can formulate, in complete analogy, that u is rationally conjugate to the group element g if and only if $\varphi(u^i) = \varphi(g^i)$ for all irreducible Brauer characters φ and integers i , or $\mu(\xi, u, \varphi) = \mu(\xi, g, \varphi)$ for all φ and ξ .

Let ψ be either an irreducible ordinary character, or an irreducible Brauer character of G relative to p when u is p -regular. The multiplicities $\mu(\xi, u, \psi)$ can be computed from the values $\psi(u^i)$ by Fourier inversion, and taking the Galois action into account, the formula reads—with ζ denoting a primitive n -th root of unity:

$$\mu(\xi, u, \psi) = \frac{1}{n} \sum_{d|n} \text{Tr}_{\mathbb{Q}(\zeta^d)/\mathbb{Q}}(\psi(u^d)\xi^{-d}).$$

Partial augmentations come into play by means of

$$(1) \quad \psi(u) = \sum_{x \in G} \varepsilon_x(u) \psi(x),$$

where it is understood that a Brauer character vanishes off the p -regular elements. This is obvious if ψ is an ordinary character, and can be seen as a consequence of the non-singularity of the Cartan matrix if ψ is a Brauer character, remembering the fact that the partial augmentations of the p -regular torsion unit u at p -singular group elements are zero. Combining both equations, one obtains

$$(2) \quad \mu(\xi, u, \psi) = \frac{1}{n} \sum_{x \in G} \sum_{d|n} \varepsilon_x(u^d) \text{Tr}_{\mathbb{Q}(\zeta^d)/\mathbb{Q}}(\psi(x)\xi^{-d}).$$

This should be seen as a linear system of equations with rational coefficients in the unknown partial augmentations. Note that the $\mu(\xi, u, \psi)$ are non-negative integers, bounded above by $\psi(1)$.

3. APPLICATION OF THE LUTHAR-PASSI METHOD

Let u be a torsion unit in $V(\mathbb{Z}A_6)$. The order of u is a divisor of the exponent of A_6 , which is $2^2 \cdot 3 \cdot 5$. The character table of A_6 is shown in Table 1 in the form obtained by requiring `CharacterTable("A6")` in GAP [10] (dots indicate zeros). So kx is the “ x ”-th class of elements of order k , and ε_{kx} will denote the partial augmentation of u with respect to this class. We will also make use of the Brauer character tables of A_6 for the primes 2 and 3, shown in Table 2 (which can be accessed from the GAP library through the `mod` command if the character table is required in the above form). We assume that $u \neq 1$, so $\varepsilon_{1a} = 0$ by the familiar Berman–Higman result. We shall show that all but one of the partial augmentations of u vanish if its order is not 6. When u is assumed to have order 6, the Luthar–Passi method is not strong enough to exclude the possibility of having $(\varepsilon_{2a}, \varepsilon_{3a}, \varepsilon_{3b})$ equal to $(-2, 1, 2)$ or $(-2, 2, 1)$. Note that we have to consider the cases when u has order 2, 4, 3, 5 (then vanishing of all but one of the partial augmentations

	1a	2a	3a	3b	4a	5a	5b
χ_{1a}	1	1	1	1	1	1	1
χ_{5a}	5	1	2	-1	-1	.	.
χ_{5b}	5	1	-1	2	-1	.	.
χ_{8a}	8	.	-1	-1	.	α_1	α_2
χ_{8b}	8	.	-1	-1	.	α_2	α_1
χ_{9a}	9	1	.	.	1	-1	-1
χ_{10a}	10	-2	1	1	.	.	.

Irrational entries: $\alpha_1 = (1 + \sqrt{5})/2 = 1 + \nu + \nu^4$ and
 $\alpha_2 = (1 - \sqrt{5})/2 = 1 + \nu^2 + \nu^3$, where $\nu = \exp(2\pi i/5)$.

TABLE 1. Character table of A_6

	1a	3a	3b	5a	5b
$\varphi_{2,1a}$	1	1	1	1	1
$\varphi_{2,4a}$	4	1	-2	-1	-1
$\varphi_{2,4b}$	4	-2	1	-1	-1
$\varphi_{2,8a}$	8	-1	-1	α_1	α_2
$\varphi_{2,8b}$	8	-1	-1	α_2	α_1

	1a	2a	4a	5a	5b
$\varphi_{3,1a}$	1	1	1	1	1
$\varphi_{3,3a}$	3	-1	1	α_1	α_2
$\varphi_{3,3b}$	3	-1	1	α_2	α_1
$\varphi_{3,4a}$	4	.	-2	-1	-1
$\varphi_{3,9b}$	9	1	1	-1	-1

TABLE 2. Brauer character tables of A_6 mod 2 and mod 3

will be shown), when u has order 10 or 15 (a contradiction must be reached), and when u has order 6 (with the final contradiction to be obtained in the next section). Remark 1 on the vanishing of certain partial augmentations will be used without further mention. Finally, we remark that A_6 has automorphisms which permute either the classes **3a** and **3b** or the classes **5a** and **5b** (leaving the remaining classes fixed), which helps from repeating steps.

A subscript $*$ will always stand as a placeholder for all possible subscripts. We denote by Θ_* a representation of G , over a sufficiently large field, which affords the Brauer character φ_* , and write $\Theta_*(u) \sim \text{diag}(\lambda_1, \dots, \lambda_{\varphi_*(1)})$ to indicate that $\Theta_*(u)$ has eigenvalues $\lambda_1, \dots, \lambda_{\varphi_*(1)}$ (multiplicities taken into account), which will, in the sense of Brauer, be identified with complex roots of unity.

When u has order 2. There is only one class of elements of order 2, so there is nothing to do.

When u has order 4. Taking augmentation gives $\varepsilon_{2a} + \varepsilon_{4a} = 1$, so $|-2\varepsilon_{2a} + 1| = |-\varepsilon_{2a} + \varepsilon_{4a}| = |\varphi_{3,3a}(u)| < 3$ (strict inequality as $|\varphi_{3,3a}(u^2)| = |\varphi_{3,3a}(2a)| \neq 3$). Thus $\varepsilon_{2a} \in \{0, 1\}$, and one of ε_{2a} and ε_{4a} is zero (of course, this must be ε_{2a}).

When u has order 3. Taking augmentation gives $\varepsilon_{3a} + \varepsilon_{3b} = 1$. Thus $|3\varepsilon_{3a} - 2| = |\varepsilon_{3a} - 2\varepsilon_{3b}| = |\varphi_{2,4a}(u)| \leq 4$ and $|-3\varepsilon_{3a} + 1| = |-2\varepsilon_{3a} + \varepsilon_{3b}| = |\varphi_{2,4b}(u)| \leq 4$. So $\varepsilon_{3a} \in \{0, 1, 2\} \cap \{-1, 0, 1\} = \{0, 1\}$, and we are done.

When u has order 5. Taking augmentation gives $\varepsilon_{5a} + \varepsilon_{5b} = 1$, so $\varphi_{3,3a}(u) = (1 + \nu + \nu^4)\varepsilon_{5a} + (1 + \nu^2 + \nu^3)\varepsilon_{5b} = 1 + (1 - \varepsilon_{5a})(\nu^2 + \nu^3) + \varepsilon_{5a}(\nu + \nu^4)$. Since $\varphi_{3,3a}(u)$ is a sum of three fifth root of unity, it follows that $\varepsilon_{5a} \in \{0, 1\}$, as desired.

By slight abuse of notation, we shall also write kx for a representative of the class kx .

When u has order 10. Taking augmentation gives $\varepsilon_{2a} + \varepsilon_{5a} + \varepsilon_{5b} = 1$. Thus $\varphi_{3,3a}(u) = -1 + \varepsilon_{5a}(3 + \sqrt{5})/2 + \varepsilon_{5b}(3 - \sqrt{5})/2$. We can assume without loss of generality that u^6 is rationally conjugate to $5a$. Also u^5 is rationally conjugate to $2a$. We have $\Theta_{3,3a}(5a) \sim \text{diag}(1, \nu, \nu^4)$ and $\Theta_{3,3a}(2a) \sim \text{diag}(1, -1, -1)$. The eigenvalues of $\Theta_{3,3a}(u)$ are products of the diagonal entries of the diagonal matrices just depicted, always taken one from each matrix, in such a way that all entries are involved. Since $\varphi_{3,3a}(u)$ is real, it follows that $\Theta_{3,3a}(u) \sim \text{diag}(1, -\nu, -\nu^4)$ and $\varphi_{3,3a}(u) = (3 - \sqrt{5})/2$. Comparing the given values for $\varphi_{3,3a}(u)$ results in an equation which has no integral solution (reduce modulo 3), a contradiction.

When u has order 15. Then we can assume without loss of generality that u^{10} is rationally conjugate to $3a$ and that u^6 is rationally conjugate to $5a$. Setting $\zeta = \exp(2\pi i/3)$, we have $\Theta_{2,4a}(3a) \sim \text{diag}(1, 1, \zeta, \zeta^2)$ and $\Theta_{2,4a}(5a) \sim \text{diag}(\nu, \nu^2, \nu^3, \nu^4)$. The eigenvalues of $\Theta_{2,4a}(u)$ are the products of the diagonal entries of these diagonal matrices, taken in a suitable order. But there is no possibility to choose an ordering such that the sum of the products becomes rational, a contradiction.

When u has order 6. Taking augmentation gives $\varepsilon_{2a} + \varepsilon_{3a} + \varepsilon_{3b} = 1$. Thus $\chi_{5a}(u) = 1 + \varepsilon_{3a} - 2\varepsilon_{3b}$ and $\chi_{5b}(u) = 1 - 2\varepsilon_{3a} + \varepsilon_{3b}$. Now, u^3 is rationally conjugate to $2a$, and we can assume without loss of generality that u^2 , and hence also u^4 , is rationally conjugate to $3b$. Write D_{5a} and D_{5b} for ordinary representations affording χ_{5a} and χ_{5b} , respectively. We have $D_{5a}(2a) \sim \text{diag}(1, 1, 1, -1, -1)$ and $D_{5a}(3b) \sim \text{diag}(1, \zeta, \zeta^2, \zeta, \zeta^2)$. The eigenvalues of $D_{5a}(u)$ are products of the diagonal entries of these diagonal matrices, suitably taken. Their sum $\chi_{5a}(u)$ is rational, which forces

$$(3) \quad D_{5a}(u) \sim \text{diag}(1, \zeta, \zeta^2, -\zeta, -\zeta^2)$$

and $\chi_{5a}(u) = 1$. Thus $\varepsilon_{3a} = 2\varepsilon_{3b}$ and $\chi_{5b}(u) = 1 - 3\varepsilon_{3b}$. We have $D_{5b}(2a) \sim \text{diag}(1, 1, 1, -1, -1)$ and $D_{5b}(3b) \sim \text{diag}(1, 1, 1, \zeta, \zeta^2)$. In the same way as before, we obtain that either

$$(4) \quad D_{5b}(u) \sim \text{diag}(1, -1, -1, \zeta, \zeta^2)$$

and $\chi_{5b}(u) = -2$, or $\chi_{5b}(u) = 4$. If $\chi_{5b}(u) = 4$, then $\varepsilon_{3b} = -1$, $\varepsilon_{3a} = -2$ and $\varepsilon_{2a} = 4$; but then $\chi_{10a}(u) = -11$, a contradiction. Hence $\chi_{5b}(u) = -2$, and $\varepsilon_{3b} = 1$, $\varepsilon_{3a} = 2$ and $\varepsilon_{2a} = -2$, one of the possibilities mentioned at the beginning.

4. NON-EXISTENCE OF A UNIT OF ORDER 6 IN $V(\mathbb{Z}A_6)$

Finally, we assume that there exists a unit u of order 6 in $V(\mathbb{Z}A_6)$, and are looking for a contradiction. According to the previous section, we can assume that u has partial augmentations $(\varepsilon_{2a}, \varepsilon_{3a}, \varepsilon_{3b})$ equal to $(-2, 2, 1)$, that u^3 is rationally conjugate to $2a$ and that u^2 is rationally conjugate to $3b$. In particular, the character values of all powers of u are known, from which the eigenvalues of representing matrices can be computed.

We shall take advantage only of representations D_{5a} and D_{5b} affording the irreducible characters χ_{5a} and χ_{5b} , for which the eigenvalues of $D_{5a}(u)$ and $D_{5b}(u)$ are already given by (3) and (4), taking into account that χ_{5a} and χ_{5b} have the same 3-modular constituents, namely the principal character and the 3-modular character of degree 4 (as can be seen from Tables 1 and 2).

Set $R = \mathbb{Z}_{(3)}$, the localization of $\mathbb{Z}$ at the prime 3, and $k = R/3R$ (the prime field of characteristic 3). For an RA_6 -lattice L , we let $\bar{L}$ stand for the kA_6 -module $L/3L$, and we also let $x \mapsto \bar{x}$ denote the natural homomorphism from RA_6 to kA_6 .

There are two doubly transitive actions of A_6 on a set with six elements, corresponding to the two conjugacy classes of subgroups of A_6 which are isomorphic to A_5 , and D_{5a} and D_{5b} can be chosen to be the associated deleted permutation representations. Denote by L_a and L_b the non-trivial factor lattices of the RA_6 -permutation lattices arising from these actions, so that a matrix representation D_{5*} affording χ_{5*} can be obtained with respect to an R -basis of L_* .

Set $e = (1 + u^3)/2$ and $f = 1 - e$, so $1 = e + f$ is an idempotent decomposition in RA_6 . Let $w = e + fu^2 \in RA_6$. From (4) one sees that $D_{5b}(w)$ is the identity matrix. In particular, $\bar{w}$ acts as the identity on the composition factors of $\bar{L}_b$, which are also the composition factors of $\bar{L}_a$, as already noted.

Now we turn our attention to the other representation. Set $L = L_a$. Note that the simple kA_6 -module $\tilde{M}$ of dimension 4 is a submodule of $\bar{L}$, namely $\tilde{M} = \bar{L}I(kA_6)$ where $I(kA_6)$ denotes the augmentation ideal of kA_6 . (If we set $M = LI(RA_6)$, then M is a sublattice of L of index 3 and $\tilde{M}$ is the image of M under the map $L \rightarrow \bar{L}$.) We have $L = Le \oplus Lf$. From (3) one sees that Lf is an R -lattice of rank 2 on which w acts nontrivially by multiplication with u^2 , a unit of order 3. But $\bar{L}f = \bar{L}\bar{f} \subseteq \tilde{M}$, so $\bar{w}$ acts trivially on $\bar{L}\bar{f}$, by the preceding paragraph. So we have obtained a contradiction, as the kernel of the natural map $\mathrm{GL}_2(R) \rightarrow \mathrm{GL}_2(k)$ has no 3-torsion. (A particular case of a well-known lemma due to Minkowski; see, for example, the beautiful discussion in [11, 5.2].)

REFERENCES

- [1] Ashwani K. Bhandari and Indar S. Luthar, *Certain conjugacy classes of units in integral group rings of metacyclic groups*, J. Number Theory **18** (1984), no. 2, 215–228.
- [2] V. Bovdi, C. Höfert, and W. Kimmerle, *On the first Zassenhaus conjecture for integral group rings*, Publ. Math. Debrecen **65** (2004), no. 3-4, 291–303.
- [3] V. A. Bovdi, E. Jespers, and A. B. Kononov, *Integral group ring of the first Janko simple group*, manuscript (eprint [arXiv:math.GR/0608441](https://arxiv.org/abs/math.GR/0608441)).
- [4] V. A. Bovdi and A. B. Kononov, *Integral group ring of the first Mathieu simple group*, Groups St. Andrews 2005, vol. 1, London Math. Soc. Lecture Note Ser., no. 340, Cambridge Univ. Press, Cambridge, 2007, pp. 237–245.
- [5] ———, *Integral group ring of the Mathieu simple group M_{23}* , Comm. Algebra, to appear (eprint [arXiv:math.RA/0612640](https://arxiv.org/abs/math.RA/0612640)).
- [6] V. A. Bovdi, A. B. Kononov, and S. Siciliano, *Integral group ring of the Mathieu simple group M_{12}* , Rend. Circ. Mat. Palermo (2), to appear (eprint [arXiv:math.RA/0612638](https://arxiv.org/abs/math.RA/0612638)).
- [7] Victor Bovdi and Martin Hertweck, *Zassenhaus conjecture for central extensions of S_5* , J. Group Theory, to appear (eprint [arXiv:math.RA/0609435](https://arxiv.org/abs/math.RA/0609435)).
- [8] James A. Cohn and Donald Livingstone, *On the structure of group algebras. I*, Canad. J. Math. **17** (1965), 583–593.
- [9] Michael A. Dokuchaev, Stanley O. Juriaans, and César Polcino Milies, *Integral group rings of Frobenius groups and the conjectures of H. J. Zassenhaus*, Comm. Algebra **25** (1997), no. 7, 2311–2325.
- [10] The GAP Group, *GAP — Groups, Algorithms, and Programming, Version 4.4*, 2006, (<http://www.gap-system.org>).

- [11] Robert M. Guralnick and Martin Lorenz, *Orders of finite groups of matrices*, Groups, Rings and Algebras, A conference in Honor of Donald S. Passman (W. Chin, J. Osterburg, and D. Quinn, eds.), Contemp. Math., vol. 420, Amer. Math. Soc., Providence, RI, 2006, pp. 141–161.
- [12] Martin Hertweck, *Torsion units in integral group rings of certain metabelian groups*, Proc. Edinburgh Math. Soc. (2), to appear.
- [13] ———, *Partial augmentations and Brauer character values of torsion units in group rings*, Comm. Algebra, to appear (eprint [arXiv:math.RA/0612429v2](https://arxiv.org/abs/math.RA/0612429v2)).
- [14] ———, *On the torsion units of some integral group rings*, Algebra Colloq. **13** (2006), no. 2, 329–348.
- [15] Christian Höfert and Wolfgang Kimmerle, *On torsion units of integral group rings of groups of small order*, Groups, rings and group rings, Lect. Notes Pure Appl. Math., vol. 248, Chapman & Hall/CRC, Boca Raton, FL, 2006, pp. 243–252.
- [16] I. S. Luthar and A. K. Bhandari, *Torsion units of integral group rings of metacyclic groups*, J. Number Theory **17** (1983), no. 2, 270–283.
- [17] I. S. Luthar and I. B. S. Passi, *Zassenhaus conjecture for A_5* , Proc. Indian Acad. Sci. Math. Sci. **99** (1989), no. 1, 1–5.
- [18] I. S. Luthar and Poonam Sehgal, *Torsion units in integral group rings of some metacyclic groups*, Res. Bull. Panjab Univ. Sci. **48** (1998), no. 1-4, 137–153 (1999).
- [19] I. S. Luthar and Poonam Trama, *Zassenhaus conjecture for certain integral group rings*, J. Indian Math. Soc. (N.S.) **55** (1990), no. 1-4, 199–212.
- [20] ———, *Zassenhaus conjecture for S_5* , Comm. Algebra **19** (1991), no. 8, 2353–2362.
- [21] Z. Marciniak, J. Ritter, S. K. Sehgal, and A. Weiss, *Torsion units in integral group rings of some metabelian groups. II*, J. Number Theory **25** (1987), no. 3, 340–352.
- [22] Mohamed Ahmed M. Salim, *Torsion units in the integral group ring of the alternating group of degree 6*, manuscript, 2006.

UNIVERSITÄT STUTTGART, FACHBEREICH MATHEMATIK, IGT, PFAFFENWALDRING 57, 70550 STUTTGART, GERMANY

E-mail address: hertweck@mathematik.uni-stuttgart.de